




Contents lists available at ScienceDirect

## Journal of Anthropological Archaeology

journal homepage: [www.elsevier.com/locate/jaa](http://www.elsevier.com/locate/jaa)

## Faunal evidence for a difference in clothing use between Neanderthals and early modern humans in Europe

Mark Collard <sup>a,b,c,\*</sup>, Lia Tarle <sup>b</sup>, Dennis Sandgathe <sup>a,b,d</sup>, Alexander Allan <sup>a,b</sup>

<sup>a</sup> Human Evolutionary Studies Program, Simon Fraser University, Burnaby, British Columbia, Canada

<sup>b</sup> Department of Archaeology, Simon Fraser University, Burnaby, British Columbia, Canada

<sup>c</sup> Department of Archaeology, University of Aberdeen, Aberdeen, Scotland, UK

<sup>d</sup> European Section, University of Pennsylvania Museum of Archaeology and Anthropology, Philadelphia, PA, USA

## ARTICLE INFO

## Article history:

Received 12 January 2016

Revision received 3 July 2016

Available online xxxx

## Keywords:

Neanderthals

Early modern humans

Oxygen Isotope Stage 3

Specialized cold weather clothing

Cape-like clothing

Mustelidae

Canidae

Mousterian

Aurignacian

Gravettian

## ABSTRACT

In this paper we report a study designed to shed light on the possibility that clothing differences played a role in the replacement of the Neanderthals by early modern humans. There is general agreement that early modern humans in Europe utilized specialized cold weather clothing, but the nature of the clothing used by Neanderthals is debated. Some researchers contend that they did not use clothes. Others argue that they were limited to cape-like clothing. Still others aver that their clothing was not substantively different in terms of thermal effectiveness from that of early modern humans. To test among these hypotheses, we employed a novel line of evidence—the bones of animals whose skins may have been made into clothing. We used an ethnographic database to identify mammalian families that were used to create cold weather clothing in the recent past. We then compared the frequency of occurrence of these families in European archaeological deposits associated with early modern humans and Neanderthals. We obtained two main results. One is that mammalian families used for cold weather clothing occur in both early modern human- and Neanderthal-associated strata. The other is that three of the families—leporids, canids, and mustelids—occur more frequently in early modern human strata than in Neanderthal strata. There is reason to believe that the greater frequency of canid and mustelid remains in early modern human strata reflects the use of garments with fur trim. Thus, these findings are most consistent with the hypothesis that Neanderthals employed only cape-like clothing while early modern humans used specialized cold weather clothing. We end by discussing the implications of this hypothesis for the debate about the replacement of the Neanderthals by early modern humans.

© 2016 Published by Elsevier Inc.

### 1. Introduction

Recently significant progress has been made in our understanding of modern human origins. For several decades there was debate about the nature of the appearance of modern humans outside of Africa. Some argued that *Homo sapiens* originated in Africa around 200,000 years ago (kya) and then spread throughout the rest of the world, replacing or absorbing regional groups of non-modern hominins as they went (Stringer, 2002). Others asserted that *H. sapiens* evolved in different regions from different groups of non-modern hominins over the course of the last two million years (Wolpoff et al., 2000). This dispute has been resolved in the last few years, as a result of new fossil discoveries and the development

of novel methods (e.g. ancient DNA) (Collard and Dembo, 2013). Today, there is widespread agreement that *H. sapiens* originated in Africa about 200 kya and migrated into the other regions of the world 100,000–150,000 years later (Cartmill and Smith, 2009; Klein, 2009). Even those researchers who were once the main proponents of the multiregional evolution model now accept that migration from Africa within the last 100,000 years played an important role in the appearance of *H. sapiens* outside of Africa (Wolpoff et al., 2004). Now that the out of Africa versus multiregional evolution debate has been resolved in favour of the former model, attention has shifted to elucidating the details of the process by which the migrating early modern humans replaced the various regional groups of non-modern hominins.

In western Eurasia, the non-modern hominins replaced by the migrating early modern humans were the Neanderthals. Neanderthals were close relatives of modern humans—so close in fact that the two species seem to have been able to interbreed (e.g.

\* Corresponding author at: Human Evolutionary Studies Program and Department of Archaeology, Simon Fraser University, 8888 University Drive, Burnaby, British Columbia V5A 1S6, Canada.

E-mail address: [mcollard@sfu.ca](mailto:mcollard@sfu.ca) (M. Collard).

Green et al., 2010; Fu et al., 2014). The Neanderthals had brains that were similar in size to those of *H. sapiens*, a long, low cranial vault with pronounced brow-ridges, and a large, prognathic face (Cartmill and Smith, 2009). Neanderthals were stocky. Their average body mass and stature have been estimated as 72 kg and 161–165 cm, respectively (Ruff et al., 1997; Feldesman et al., 1990). They also had relatively short forearms and lower legs (Ruff, 1993; Holliday, 1997). The size and shape of the Neanderthal body are widely accepted to be adaptations to glacial conditions (Ruff, 1993; Holliday, 1997). Neanderthals lived in small, dispersed groups, and specialized in hunting large game (Stiner, 2001; Stiner et al., 2009). They made sophisticated stone tools, but evidence from several sites indicates that their use of fire was restricted to warm periods, which suggests they may not have been able to create fire at will but only take advantage of naturally occurring fires (Sandgathe et al., 2011). Additionally, they did not build structures or utilize symbols on a regular basis (Klein, 2003). Genetic and morphological data suggest Neanderthals were a distinct species by at least 200 kya (Cartmill and Smith, 2009). Early modern humans joined Neanderthals in Europe ca. 42 kya, during Oxygen Isotope Stage 3 (OIS3) (Stringer, 2006). Within a few thousand years, the Neanderthals had disappeared. Current evidence indicates they went extinct about 41–39 kya (Higham et al., 2014). With regard to geographic range, the Neanderthals were a western Eurasian species. Their fossilized remains have been found from Wales in the north to Israel in the south, and from Portugal in the west to Central Asia in the east (Klein, 2003; Krause et al., 2007). So far, no Neanderthal fossils have been found in Africa, South Asia, or East Asia.

Why early modern humans were able to replace Neanderthals is contested. Some researchers argue that early modern humans out-competed Neanderthals because they were able to exploit more resilient and reliable resources, such as rabbits, fish, and plants that require processing to eat (Stiner, 2001; Mellars, 2004; Stiner and Kuhn, 2006; Richards and Trinkaus, 2009). Others aver that Neanderthals did not disappear as a consequence of competition with early modern humans. According to Stewart (2007), for example, the fact that Neanderthals died out in Europe at the same time as two “interglacial survivors,” the straight-tusked elephant (*Elephas antiquus*) and Merck’s rhino (*Stephanorhinus kirchbergensis*), indicates that, contrary to the current consensus, Neanderthals were warm adapted rather than cold adapted. The corollary of this, Stewart contends, is that Neanderthals went extinct because they were unable to cope with the substantial decrease in temperature associated with OIS3. Finlayson (2004, 2009) and Jiménez-Espejo et al. (2007) have also argued that the Neanderthals were driven to extinction by the effects of climate change rather than competition with early modern humans. Still others have proposed that a combination of competition with early modern humans and the effects of climate change were responsible for the Neanderthals’ disappearance (Stringer et al., 2003). These researchers suggest that during OIS3, rapid climatic fluctuations destabilized the environment, and the combined stress of an unstable resource base and the arrival of new competitors drove the Neanderthals to extinction.

The study reported here was designed to shed light on the possibility that differences in clothing played a role in the replacement of Neanderthals by early modern humans. Jim O’Connell was, we believe, the first person to suggest such might be the case. Jim put forward this idea in his 2006 contribution to the modern human origins debate, “How did modern humans displace Neanderthals? Insights from hunter-gatherer ethnography and archaeology” (O’Connell, 2006). Jim’s thesis in this paper was that the replacement of the Neanderthals by early modern humans may have been an episode of competitive exclusion in which differences in diet breadth were crucial. He argued that early modern humans

had a broader diet than the Neanderthals, and went on to suggest that this would have allowed them to occupy a wider array of habitats than the Neanderthals. As a result of this, he continued, early modern humans eventually would have driven the Neanderthals from their former range. While discussing the archaeological evidence that supports this scenario, Jim highlighted a critical prerequisite for early modern humans to have been able to occupy a wider range of habitats than Neanderthals—namely that they would have had to invest more heavily in technologies for coping with cold conditions (e.g. hearths, shelter, and clothing) than Neanderthals. The idea that clothing played an important role in the replacement of the Neanderthals by early modern humans was subsequently elaborated by Gilligan (2007) and Wales (2012).

The impact of differences in Neanderthal and early modern human clothing could have been substantial. As is well known, prolonged exposure to cold in the absence of adequate clothing can lead to frostbite and hypothermia, and eventually, death. At the extreme, then, differences in clothing could have had an impact on the health and perhaps even the survivorship of Neanderthals compared to early modern humans. Even if this were not the case, the impact of clothing differences could still have been considerable. For example, given the need to avoid frostbite and hypothermia, such differences could have influenced the length of the daily “time window” for foraging, and limited the latitude and elevation at which foraging was possible, which in turn could have affected daily foraging return rates. Differences in clothing may have affected foraging in other ways too. Ethnographic work indicates that insufficiently warm clothing can hinder hunting tactics involving long periods of inactivity, such as ambush hunting (Stenton, 1991). Therefore, clothing differences could have impacted the effectiveness of foraging, resulting in a difference in daily foraging return rates. This in turn could have led to a difference in calorie intake and, ultimately, inter-birth interval (Froehle and Churchill, 2009). Thus, even if differences in clothing did not affect health and survivorship directly, they could have played a role in the replacement event via their impact on reproductive rate and demography.

Currently, it is unclear whether there was a difference in early modern human and Neanderthal clothing. There is general agreement that as early modern humans moved into glacial Europe, they would have adopted highly insulative specialized cold weather clothing, involving multiple fitted garments made from well-tanned, pliable hides.<sup>1</sup> This is based, in part, on the recovery of bone needles at early modern human sites in Africa and Eurasia (Backwell et al., 2008; Hoffecker, 2005a). In Africa, a bone needle-like implement has been recovered from deposits dating to ca. 61 kya at the site of Sibudu, South Africa (Backwell et al., 2008). The oldest eyed bone needles from Eurasia date to 37–40 kya and are generally accepted to be associated with modern humans (Golovanova et al., 2010a, 2010b). There is also evidence that early modern humans regularly processed hides. Ethnographic and archaeological data indicate that lithic endscrapers are specialized tools for intensive hide scraping (e.g. Hayden, 1990; Jefferies, 1990; Shott and Weedman,

<sup>1</sup> At the moment, there is no generally accepted terminology for discussing craft-produced clothing. For example, Hayden (1990) suggested a tripartite classification of “basic capes,” “improved capes,” and “luxury garments,” while Gilligan (2007) distinguished between “simple clothing” and “complex clothing.” We are not content with either of these schemes. Accordingly, we have elected to use the terms “cape-like clothing” and “specialized cold weather clothing” to refer to the two types of ensemble that have so far featured in the debate about Neanderthal and early modern human clothing use. We believe the meaning of “cape-like clothing” should be self-evident. The term “cold weather clothing” is often used by researchers who work on clothing performance to refer to ensembles of garments that are designed to protect against extremely cold environments such as the Arctic and high mountains (e.g. Oakes et al., 1995). We added “specialized” to make it even clearer that the garments are specifically designed for cold weather.

2007; Loebel, 2013), and endscrapers are common in early modern human sites in Eurasia. Additional support for the hypothesis that early modern humans used specialized cold weather clothing is provided by ca. 24,000 year old carved ivory figurines from Russia that appear to depict individuals wearing fur parkas (Hoffecker, 2005a).

In contrast, the nature of the clothing used by Neanderthals is debated. Some researchers have suggested Neanderthals did not use clothing. Kittler et al. (2003), for example, used molecular techniques to investigate the divergence between human head and body lice. They reasoned that this divergence would have occurred after hominins began to wear clothes regularly and therefore dating it should provide a *terminus post quem* for the origin of clothing. They also reasoned that the genetic diversity of body lice from different regions should be informative about the origin of the use of clothing. Kittler et al. found that head and body lice diverged about 72,000 ± 42,000 years ago, which postdates the first appearance of early modern humans. They also found much greater genetic diversity among African body lice than among body lice from other regions. Kittler et al. concluded from these findings that the use of clothing likely originated with early modern humans in Africa, and that archaic hominins like the Neanderthals probably did not use clothing. The following year Kittler et al. (2004) reported a revised date of 107 kya for the origin of body lice. Kittler et al.'s (2003, 2004) basic finding—that clothing use first developed among early modern humans living in Africa—was subsequently replicated using more sophisticated techniques by Toups et al. (2011).

Other researchers have argued that Neanderthals used clothing but did not employ garments of the same thermal effectiveness as early modern humans (Hayden, 1990, 1993; Gilligan, 2007; Wales, 2012). Drawing on ethnographic data, Hayden (1990, 1993) argued that specialized cold weather clothing was not a requirement among many historic groups living in temperate regions. Rather, it was mainly a status marker and its production depended on resource abundance. Because he could identify no evidence for true resource abundance in the Middle Palaeolithic, he suggested that Neanderthals likely used only cape-like clothing. Subsequently, Gilligan (2007) argued that the Neanderthals probably relied on cape-like clothing rather than specialized cold weather clothing because no needles and few precision cutting tools have been recovered from Neanderthal sites. Wales (2012) has also argued that Neanderthal clothing would have been less thermally effective than that of early modern humans. He modeled climate conditions at early modern human and Neanderthal sites in Europe, and found that early modern human sites were frequently in areas that would have required them to cover 80–90% of their bodies, whereas Neanderthal sites were located in areas that would have required only 70–80% of the body to be covered. Wales concluded from this that while early modern humans would often have needed to employ specialized cold weather clothing, Neanderthals would rarely have needed to do more than drape themselves with the fur of a large-bodied animal.

Still other researchers have suggested that the clothing of the Neanderthals would have been similar in terms of thermal effectiveness to the clothing used by early modern humans. White (2006) is one such researcher. Focusing on the situation in Britain, he argued that the low temperatures during OIS3 would have required Neanderthals to wear tailored fur clothing. He also argued that cape-like clothing would not have been an option because it would have impeded arm movements. Sørensen (2009) is another proponent of the hypothesis that Neanderthal clothing would have been similar to that of early modern humans. He modeled Neanderthal energetics with different amounts of clothing, and concluded that Neanderthals living in northern Europe would have had to cover all but small portions of their bodies with fitted clothing as early as 125 kya, long before the start of OIS3. According to

Sørensen, the absence of bone needles at Neanderthal sites does not mean they only employed cape-like clothing. He suggested that Neanderthals would have been able to make fitted clothing using what he describes as “awl-like points” and “knife-like blades” to cut strips of skin and join skins together (p. 2203).

In our study, we focused on a line of evidence that so far has received relatively little attention in the debate about the possibility of a difference in the clothing of early modern humans and Neanderthals—the bones of animals whose skins may have been used to produce clothing (see White [2006] for a rare exception). The study had three parts. First, we identified mammalian families that were present in Europe during OIS3. This was accomplished with the aid of the Stage 3 Project Faunal Database, which records the presence of mammal species at 493 European archaeological and palaeontological sites dated to 60–20 kya (Stewart et al., 2003). Next, we identified which of the Stage 3 mammalian families were used to manufacture cold weather clothing by recent mid-to-high latitude small-scale societies. This was achieved with the assistance of a large ethnographic database. In the last part of the study, we returned to the Stage 3 Project Faunal Database. We targeted families that the second part of the study indicated were used in cold weather clothing, and tested for significant differences in the frequencies of their remains in Neanderthal- and early modern human-associated strata. We focused on the 333 strata in the Stage 3 Faunal Database that contain Mousterian, Aurignacian, or Gravettian artefacts. Neanderthals are widely accepted to have manufactured the Mousterian, while early modern humans are generally thought to have produced the Aurignacian and Gravettian (e.g. van Andel and Davies, 2003; O'Connell, 2006). The Mousterian, which is often treated as synonymous with the Middle Palaeolithic period, emerged ca. 250 kya and disappeared around 41–39 kya (Adler et al., 2014; Higham et al., 2014; Richter, 2011). The Aurignacian appeared around 42 kya and is the first industry in Europe uniquely associated with early modern humans (Douka et al., 2014). The Gravettian overlapped with the end of the Aurignacian and lasted until around 24 kya (Hoffecker, 2005b). Thus, together, the Aurignacian and Gravettian cover the period of coexistence between Neanderthals and early modern humans.

## 2. Materials and methods

To begin with, we identified mammalian taxa that were present in Europe during OIS3, using the Stage 3 Project Faunal Database (<http://www.esc.cam.ac.uk/research/research-groups/research-projects/stage-three-project/stage-three-project-database-downloads>). The Stage 3 Project Faunal Database contains data from most of the archaeological sites from Europe that have yielded mammalian faunal remains and been radiometrically dated to 60–20 kya, as well as a number of European palaeontological sites that fall in the same time range (Stewart et al., 2003). OIS3 began ca. 60 kya and ended ca. 24 kya. However, the members of the Stage 3 Project elected to include sites up to and including 20 kya in the database in order to incorporate the start of the LGM, which began ca. 21 kya (van Andel, 2003). For the purposes of the Stage 3 Project, the geographic extent of Europe was defined as the western Palaeartic, bordered by the Atlantic to the West, the Mediterranean to the South, the Arctic Ocean to the North, and the 40°E longitude line to the East (van Andel and Davies, 2003). Mammalian taxa are recorded as present/absent in the strata included in the database.

In the second part of the study, we searched the electronic version of the eHRAF World Cultures Database (<http://hraf.yale.edu/online-databases/ehraf-world-cultures/>) for mentions of the use of the Stage 3 mammals in cold weather clothing produced by recent non-industrial groups from mid-to-high latitudes. The

eHRAF World Cultures Database is a cross-cultural database that contains information on numerous aspects of the cultural and social lives of a worldwide sample of 258 ethnographic groups. In searching the eHRAF World Cultures Database, common names were used rather than formal taxonomic names. The common names were taken from *Walker's Mammals of the World* (Nowak, 1999). We focused on the family level because the common names of many taxa are not unique to a single genus or species (e.g. “fox” can refer to the *Vulpes* genus or the *Alopex* genus), and because several taxa that have been recovered from archaeological strata that date to OIS3 are now extinct and therefore could not have been utilized by recent human groups. Focusing on the family level allowed extinct taxa to be included in the analysis, because extinct species and genera could be categorized as members of extant families that are used for clothing by recent groups. Because the goal of the analyses was to identify taxa used in traditional, craft-produced cold weather clothing, we did not include groups such as African Americans, Arab Americans and Cuban Americans (deemed “Regional and Ethnic Cultures” by the eHRAF World Cultures Database) or results for clothing suspected to have been mass-produced. Disregarding “Regional and Ethnic Cultures” left a total of 237 groups. Of these 237 groups, 133 come from mid-to-high latitudes, and 104 come from tropical latitudes. We defined “mid-to-high” latitude groups as those whose territories are north of or encompass the Tropic of Cancer, or south of or encompass the Tropic of Capricorn. The geographic territories of the groups were taken from the Human Relations Area Files, and latitudes from Stanford (2003). Any item of clothing described as warm or for winter or cold weather use was deemed to be “cold weather clothing.” Repeated mentions of the use of a taxon for a particular type of clothing by a single group were not counted. For example, if multiple ethnographies noted that the Chukchee used reindeer hides to make winter parkas, only one use was counted.

Lastly, we returned to the Stage 3 Project Faunal Database to test for differences between Neanderthals and early modern humans in their exploitation of fauna suitable for producing cold weather clothing. Specifically, we tested for differences in the frequency of the remains of mammalian families that the second step of the study indicated were used to produce cold weather clothing in ethnographic contexts. We compared early modern human-associated Aurignacian/Gravettian strata with Neanderthal-associated Mousterian strata, focusing on families that were often used for cold weather clothing in the ethnographic sample. We defined “often” as  $\geq 1\%$  of ethnographic uses. We did not include so-called “transitional” industries such as the Chatelperronian because of the ongoing debate about their validity as cultural entities and their association with Neanderthals versus early modern humans (e.g. Bar-Yosef and Bordes, 2010).

In selecting strata, we limited the date range to 60–20 uncal kya, so that outliers were not included. Only strata that contain faunal data and were clearly linked with the Mousterian, Aurignacian, or Gravettian were included. Potentially mixed strata were not included. Duplicate entries for a stratum were consolidated. If strata names given by the original monograph and the database’s editorial notes were contradictory, the stratum name from the original monograph was used. If the editorial notes gave additional stratigraphic information that did not contradict the original monograph, the editorial notes were used to distinguish strata. If strata were not named in the database but were associated with different industries (e.g. one unnamed Aurignacian stratum and one unnamed Mousterian stratum from the same site), both strata were included. Unnamed strata associated with the same industry at a single site that were only differentiated by elevation were consolidated. If an entry indicated that it represented more than one stratum (e.g. “stratum 10–11”), and the strata in question were already represented in the dataset, the multi-

strata data were consolidated with the existing strata (e.g. “stratum 10–11” was consolidated with “stratum 10” and “stratum 11”). However, if one of the strata was not already represented, the multi-strata data were kept as a distinct data point (e.g. if “stratum 2–9” was present as well as “strata 5” and “strata 6,” then “stratum 2–9” was kept as a distinct stratum). At some sites, only descriptions were given as strata names. In these cases, strata with similar names were consolidated. As a rule, the strata nomenclature listed in the database was trusted, unless strata names obviously did not in fact refer to strata. For example, at Sunghir, a number of burials were listed as strata, but dating reports (Dobrovolskaya et al., 2012) state that the burials come from the main Gravettian occupation. So, faunal data from the “burial layers” were consolidated with the Gravettian stratum data. Of the 333 strata that were included in the analysis, 96 were Mousterian, 132 Aurignacian, and 105 Gravettian.

To test for differences in the frequency of the remains of cold weather clothing families between Aurignacian/Gravettian strata and Mousterian strata, we used the chi-squared test. Because we effectively conducted multiple unplanned tests, we used Benjamini and Hochberg’s (1995) method of significance-level correction.

### 3. Results

The remains of members of 24 mammalian families are recorded in the Stage 3 Project Faunal Database. These families are listed in Appendix A.

Table 1 summarizes the results of our search of the eHRAF World Cultures Database. Details of animals used to make cold weather clothing were available for 77 of the 133 mid-to-high latitude societies included in the search. In total, there were 238 distinct cold weather clothing uses of the mammalian families represented in the Stage 3 Project Faunal Database. Cervidae was the most commonly utilized family; members of the deer family accounted for 32% of the cold weather clothing uses. After Cervidae, the next most commonly used families were Bovidae (16%), Mustelidae (14%), Leporidae (10%), Canidae (10%), Ursidae (6%), Sciuridae (4%), Castoridae (4%), and Felidae (2%). None of the other 15 families comprised more than 1% of cold weather clothing uses. Hereinafter, we will refer to the nine families that comprised 2% or more of the uses as “cold weather clothing families.”

The remains of cold weather clothing families occur in both Mousterian strata and Aurignacian/Gravettian strata (Table 2). However, the frequencies differ, and for some families the difference is statistically significant. When the numbers of Mousterian and Aurignacian/Gravettian strata containing the remains of the nine cold weather clothing families are compared with the chi-squared test, three families are significantly more frequent in Aurignacian/Gravettian strata than in Mousterian strata: Canidae ( $p = 0.000$ ), Leporidae ( $p = 0.000$ ), and Mustelidae ( $p = 0.000$ ) (Table 2). In contrast, none of the cold weather clothing families is significantly more frequent in Mousterian strata than in Aurignacian/Gravettian strata. Thus, cold weather clothing families occur in both early modern human-associated strata and Neanderthal-associated strata, but some families are found more frequently in the former than in the latter.

### 4. Discussion

Potentially, these results have important implications for the debate about clothing use by Neanderthals and early modern humans, but there are some complications that need to be evaluated before the implications are clear. One is that the Stage 3 Faunal Database only records the presence/absence of taxa rather than

**Table 1**

Frequency of use of mammalian families for cold-weather clothing by recent mid-to-high latitude small-scale groups. Data from eHRAF. Only families that make up  $\geq 1\%$  of uses are listed. Percentages are rounded. See text for further details.

Family	#	%
Cervidae	77	32
Bovidae	39	16
Mustelidae	34	14
Leporidae	24	10
Canidae	24	10
Ursidae	14	6
Sciuridae	10	4
Castoridae	10	4
Felidae	5	2
Other families	1	0
Total	238	

**Table 2**

Results of chi-squared tests comparing frequencies of remains of cold-weather clothing families in Mousterian and Aurignacian/Gravettian strata. Data from the Stage 3 Project Mammalian Fauna Database. P-values based on the chi-squared test. Because we effectively conducted multiple unplanned tests, we used [Benjamini and Hochberg's \(1995\)](#) method of significance-level correction. Corrected significance level = 0.017. \*\*\* = P-value significant at 0.001.

Family	% Moust	% Au/Gr	p-value
Bovidae	91	84	0.096
Canidae	51	78	0.000***
Castoridae	7	9	0.559
Cervidae	92	95	0.188
Felidae	44	40	0.492
Leporidae	23	55	0.000***
Mustelidae	20	42	0.000***
Sciuridae	18	18	0.998
Ursidae	54	45	0.136

the actual number of specimens recovered per taxon. This is potentially problematic because it means that one specimen of a taxon in a stratum was given the same weight as, say, 100 specimens of the same taxon in another stratum, in the analyses. Thus, in principle, the analyses could have obscured a numerical dominance of cold weather clothing specimens in Mousterian strata. It is surprisingly difficult to evaluate this possibility. Values for Number of Individual Specimens (NISP) and Minimum Number of Individuals (MNI) have been published for remarkably few Mousterian, Aurignacian, and Gravettian sites. However, data for a small but balanced sample of sites with published NISP values (15 Mousterian and 15 Aurignacian/Gravettian sites) suggest that this concern can be discounted. Canids, leporids, and mustelids consistently represent a higher percentage of the NISPs at the Aurignacian/Gravettian sites than at the Mousterian sites ([Table 3](#)).

A second problematic issue is that certain Late Pleistocene mammals (e.g. bears, hyenas) can be expected to have utilized some of the same caves and rock shelters as Neanderthals and early modern humans, and therefore some of the bones found in the focal strata are likely to have been introduced as a result of the activities of such species. It is possible, therefore, that the difference in the representation of cold weather clothing families in Mousterian strata and Aurignacian/Gravettian strata has nothing to do with the actions of Neanderthals and early modern humans. However, it is unlikely that the activities of cave-using non-hominin species explain the higher frequency of canid, leporid, and mustelid remains in Aurignacian/Gravettian strata. Not only is the proportion of Aurignacian/Gra these taxa vettian strata that are from caves and rock shelters lower than the proportion of Mousterian strata that are from such sites ([Table 4](#)), but also the proportion of Aurignacian/Gravettian cave/rock shelter strata that contain canids, leporids,

and mustelids is lower than the proportion of Mousterian cave/rock shelter strata that contain these taxa ([Table 5](#)).

Differential availability of species is a third concern. In principle, the difference in the representation of cold weather clothing families in Mousterian strata and Aurignacian/Gravettian strata could be due to early modern humans having greater access to members of those families rather than because they targeted them more often. This also seems unlikely. The three key families—canids, leporids, and mustelids—were ubiquitous in Europe in the recent past ([Nowak, 1999](#)) and there is no obvious reason to believe the situation would have been different in the Late Pleistocene. Temporal proximity to the Last Glacial Maximum does not explain the higher frequency of canids, leporids, and mustelids in early modern human strata either. To investigate this possibility, we removed the Gravettian strata from the dataset and re-ran the comparison. The results did not change. Canid, leporid, and mustelid remains were significantly more frequent in Aurignacian strata than in Mousterian strata ([Table 6](#)). This indicates that the difference in behaviour between Neanderthals and early modern humans was present in the earliest phase of early modern human colonization of Europe and does not reflect a greater availability of cold weather clothing families closer to the Last Glacial Maximum.

The final concern is that the pattern is real but is a consequence of differences in diet rather than differences in clothing production and use, i.e. canids, leporids, and mustelids do occur more frequently in Aurignacian/Gravettian strata than in Mousterian strata because early modern humans were hunting them more often than were Neanderthals, but early modern humans were hunting them for food rather than for the production of cold weather clothing. This possibility is more difficult to evaluate, not just because taxa can be hunted for both meat and fur, but also because humans vary so widely in terms of the items they are willing to eat. Nevertheless, we think it is unlikely that the difference in the representation of cold weather clothing families in Mousterian and Aurignacian/Gravettian strata is solely due to differences in dietary preferences. The higher frequency of Leporidae is probably due in part to early modern humans hunting them for food, but that is unlikely to be the case for Mustelidae and Canidae. A search of the eHRAF World Cultures Database indicated that no canid or mustelid genus was recorded as being used for food more than 11 times, which suggests that mustelids and canids are rarely eaten by humans living at mid to high latitudes. The corollary of this is that the greater frequency of mustelids and canids in Aurignacian/Gravettian strata is unlikely to be a consequence of these families having been hunted for food more often by early modern humans than by Neanderthals. Instead, the difference seems likely to be a consequence of a greater use of mustelid and canid pelts for cold weather clothing by early modern humans.

Given that our results appear robust, we can now consider their implications for the debate about clothing use by Neanderthals and early modern humans. To reiterate, there is general agreement that early modern humans in Europe utilized specialized cold weather clothing, but the nature of the clothing used by Neanderthals is disputed. Some researchers contend that they did not use clothes (e.g. [Kittler et al., 2003](#); [Toups et al., 2011](#)). Others argue that they were limited to ineffective, cape-like clothing ([Gilligan, 2007](#); [Wales, 2012](#)). Still others aver that their clothing was not substantively different in terms of thermal effectiveness from that of early modern humans ([Sørensen, 2009](#); [White, 2006](#)). Thus, currently there are three hypotheses with respect to differences in clothing use between Neanderthals and early modern humans: (1) the Neanderthals did not use clothing at all, while early modern humans employed specialized cold weather clothing; (2) the Neanderthals employed only cape-like clothing, which was of limited thermal effectiveness, while early modern humans used specialized cold

**Table 3**  
Comparison of Number of Individual Specimens (NISP) for canids, leporids, and mustelids at Mousterian sites vs Aurignacian/Gravettian sites. All = All non-microfauna. % = Total NISP of focal taxon as a percentage of total non-microfauna NISP.

Site	Country	Canid	Leporid	Mustelid	All	Source of data
<i>Mousterian sites</i>						
Lynford	UK	7	0	0	1363	Schreve (2006)
Geissenklosterle	Germany	45	8	0	764	Kitagawa et al. (2012)
Hohle Fels	Germany	9	1	0	659	Kitagawa et al. (2012)
Hohlenstein-Stadel	Germany	401	4	0	7245	Kitagawa et al. (2012)
Bockstein	Germany	58	7	0	1208	Kitagawa et al. (2012)
Vogelherd	Germany	8	2	0	543	Kitagawa et al. (2012)
Saint-Cesaire	France	10	3	0	1544	Morin (2012)
Pech-de-l'Azé IV	France	3	0	0	1003	Laquay (1981); Niven (2013)
Roc de Marsal	France	12	0	0	2019	Castel et al. (in press)
La Quina	France	16	0	1	5396	Debénath and Jelinek (1998)
Mujina Pecina	Croatia	5	22	4	272	Miracle (2005)
Frechet	France	19	9	1	810	Mourre et al. (2008)
Pech-de-l'Azé I	France	2	2	2	813	Soressi et al. (2008)
Grotte Tournal	France	31	34	0	1644	Boyle (1998)
Jonzac	France	0	0	0	788	Niven et al. (2012)
Total		626	92	8	26,071	
%		2.40	0.35	0.03		
<i>Aurignacian/Gravettian sites</i>						
Abri Pataud	France	82	5	0	4040	Bouchud (1975)
Roc de Combe	France	51	20	22	1694	Soulier and Mallye (2012)
Saint-Cesaire	France	53	5	6	5038	Morin (2012)
La Solutre	France	25	35	0	3586	Bemilli and Bayles (2009)
Geissenklosterle	Germany	373	449	1	7046	Kitagawa et al. (2012)
Hohle Fels	Germany	166	526	1	5163	Kitagawa et al. (2012)
Hohlenstein-Stadel	Germany	254	22	0	2423	Kitagawa et al. (2012)
Bockstein	Germany	28	19	0	366	Kitagawa et al. (2012)
Vogelherd	Germany	58	27	0	4617	Kitagawa et al. (2012)
Abri Castanet	France	36	30	0	1384	Castel (2011)
Buran Kaya III	Ukraine	620	290	10	2156	Crépin et al. (2014)
Milovice G	Czech Republic	18	28	4	1326	Svoboda et al. (2005)
Willendorf I	Austria	71	9	0	224	Musil (2003)
Willendorf II	Austria	104	4	2	294	Musil (2003)
Maisieres-Canal	Belgium	64	151	1	352	Gautier (1979)
Total		2003	1620	47	39,709	
%		5.04	4.08	0.12		

**Table 4**  
Frequencies of Mousterian and Aurignacian/Gravettian strata from caves/rock shelters versus open-air sites.

	Mousterian		Aurignacian/Gravettian	
	#	%	#	%
Caves/rock shelters	83	86	174	73
Open-air sites	13	14	63	27
Total	96		237	

**Table 5**  
Frequencies of strata containing potential cave-dwelling taxa from caves/rock shelters versus open-air sites. Potential cave-dwelling taxa = leporids, canids, mustelids, ursids, sciurids, and felids. Covered = cave/rock shelter strata. Open = open-air strata.

	Mousterian		Aurignacian/Gravettian	
	Covered	Open	Covered	Open
All	63 (90%)	7 (10%)	151 (72%)	60 (28%)
Leporids	17 (77%)	5 (23%)	82 (63%)	48 (37%)
Mustelids	14 (74%)	5 (26%)	54 (55%)	45 (45%)
Canids	44 (90%)	5 (10%)	127 (68%)	59 (32%)

weather clothing; and (3) the Neanderthals used specialized cold weather clothing just like early modern humans.

Our first finding—that the remains of cold weather clothing families occur in both early modern human-associated strata and Neanderthal-associated strata—is not consistent with all of the

**Table 6**  
Results of chi-squared tests comparing frequencies of remains of cold-weather clothing families in Mousterian and Aurignacian strata. P-values based on the chi-squared test. Because we effectively conducted multiple unplanned tests, we used Benjamini and Hochberg's (1995) method of significance-level correction. Corrected significance level = 0.017. \*\*\* = P-value significant at 0.001.

Family	% Mousterian	% Aurignacian	p-value
Bovidae	91	87	0.411
Canidae	51	82	0.000***
Castoridae	7	3	0.242 <sup>a</sup>
Cervidae	92	97	0.077
Felidae	44	38	0.372
Leporidae	23	42	0.003***
Mustelidae	20	40	0.001***
Sciuridae	18	25	0.189
Ursidae	54	48	0.397

<sup>a</sup> Yates' continuity correction was applied for expected values of <5.

hypotheses. Given that, as we explained earlier, we can be reasonably confident that the canid and mustelid remains are more likely to be indicators of clothing use than they are of dietary behaviour, an implication of their occurrence in both early modern human strata and Neanderthal strata (Table 2) is that both Neanderthals and early modern humans used clothing. This conflicts with the hypothesis that Neanderthals did not use clothes at all, while early modern humans employed specialized cold weather clothing. In contrast, it is in line with the hypothesis that the Neanderthals employed only cape-like clothing while early modern humans used specialized cold weather clothing. It is also consistent with

the hypothesis that the Neanderthals used specialized cold weather clothing just like early modern humans.

Our second finding—that the remains of leporids, mustelids, and canids occur significantly more often in early modern human-associated strata than in Neanderthal-associated strata—is easier to reconcile with one of the remaining hypotheses than with the other. The higher frequency of leporids, mustelids, and canids in early modern human strata is difficult to square with the hypothesis that the Neanderthals used specialized cold weather clothing just like early modern humans. It might have been possible if some of the other cold weather clothing families occurred more frequently in Neanderthal strata than in early modern human strata. If that had been the case, it might have been possible to argue that the skins of the cold weather clothing families in question do not differ substantively in terms of thermal effectiveness from those of leporids, canids, and mustelids, and that Neanderthals and early modern humans simply chose to use different species to produce their cold weather clothing. However, given that none of the cold weather families is more common in Neanderthal-associated strata than in early modern human-associated strata, the “adaptively neutral choice” explanation is not viable. Instead, the higher frequency of leporids, mustelids, and canids in early modern human strata suggests that the clothing of early modern humans was different from that of the Neanderthals and involved the use of more fur, which would have made it more thermally effective. Thus, our results run counter to the idea that the Neanderthals used specialized cold weather clothing just like early modern humans.

By way of contrast, the higher frequency of leporids, mustelids, and canids in early modern human strata than in Neanderthal strata is consistent with the hypothesis that the Neanderthals employed only cape-like clothing while early modern humans used specialized cold weather clothing. All we need to assume is that the early modern humans’ clothing required more skins of leporids, mustelids, and canids than did the clothing of the Neanderthals. This assumption is supported by the ethnographic record, which indicates that fur trim was a key feature of the specialized cold weather clothing developed by a number of groups living in the Arctic, and that mustelid and canid pelts were favoured for such trim (Bogoraz-Tan, 1909; Pryde, 1972; Oakes et al., 1995; Cotel et al., 2004). Research carried out to shed light on this preference suggests that mustelid and canid fur makes excellent trim because its mixture of long and short hairs slows the velocity of air in the “boundary layer” at the edges of clothing, where skin is exposed, and therefore reduces heat loss (Oakes et al., 1995; Cotel et al., 2004). In addition, mustelid and canid fur sheds ice and frost easily (Oakes et al., 1995; Cotel et al., 2004). These observations suggest that the higher frequency of mustelid and canid remains in early modern human strata could well be a consequence of early modern humans adding fur trim to certain items of clothing to make them more thermally effective.

To shed further light on this possibility, we compared the number of wolverine (*Gulo gulo*) specimens in the Aurignacian/Gravettian and Mousterian strata represented in the Stage 3 Project Faunal Database. We focused on wolverines for three reasons. First, wolverine fur was widely used as ruffs on parkas and other garments by recent sub-Arctic and Arctic groups (Oakes et al., 1995). Second, work on the protection against adverse weather conditions offered by different furs indicates that wolverine fur is the best natural fur to use as a parka ruff (Oakes et al., 1995; Cotel et al., 2004; Mizell et al., 1956). It provides excellent protection from the wind, sheds hoarfrost particularly well, and is extremely durable (Oakes et al., 1995; Cotel et al., 2004; Itkonen, 1962). Lastly, contemporary trapping organizations advise that wolverine pelts are ill suited for the construction of the bodies and arms of coats because the fur is too long and the leather too heavy (e.g. <http://www.montanatrappers.org/furbearers/wolverine.htm#>). Together, these observations

suggest that the presence of wolverine in an archaeological deposit indicates the production and use of garments with fur trim.

We found a substantial disparity in the number of wolverine remains in Aurignacian/Gravettian and Mousterian strata. Fifty-six of the 237 Aurignacian/Gravettian strata contain wolverine remains, but there is not a single wolverine specimen in any of the 96 Mousterian strata. Given that there is reason to think wolverine remains indicate the use of parkas with fur ruffs, the presence of wolverine bones in numerous Aurignacian/Gravettian strata and their complete absence in Mousterian strata supports the idea that early modern humans added fur trim to their clothing to make them more thermally effective, while Neanderthals did not. This is consistent with the hypothesis that the Neanderthals employed only cape-like clothing while early modern humans used specialized cold weather clothing.

Thus, our study identifies another line of evidence that supports the hypothesis that Neanderthals used cape-like clothing while early modern humans used specialized cold weather clothing. The hypothesis is now supported by modeling work that indicates Neanderthals could not have survived during the cold periods of OIS3 without some form of clothing (Aiello and Wheeler, 2003; Wales, 2012); by the occurrence of bone needles in early modern human-associated strata but not in Neanderthal-associated strata (Hoffecker, 2005a; Golovanova et al., 2010a, 2010b); and by differences in the frequency of the remains of mammals that are known to be useful for making cold weather clothing in Neanderthal-associated and early modern human-associated strata. None of these lines of evidence is conclusive, but we believe that collectively they make a good case for favouring the hypothesis that Neanderthals used cape-like clothing while early modern humans used specialized cold weather clothing over its competitors.

There may in fact be another line of evidence that supports the hypothesis that Neanderthals used cape-like clothing while early modern humans used specialized cold weather clothing. The thermal effectiveness of hide clothing is affected by the nature and degree of processing of the hide. Pliable hide is much warmer than stiff hide, because it can be made to conform to the body. Thus, evidence for hide processing in Neanderthal strata versus early modern human strata is potentially informative about the thermal effectiveness of their clothing. There is a clear difference between Neanderthals and early modern humans with regard to the best-known hide processing tool—the endscraper. As we explained earlier, there is good evidence that endscrapers were specialized hide processing tools (e.g. Hayden, 1990; Jefferies, 1990; Shott and Weedman, 2007; Loebel, 2013). Endscrapers typically occur in high frequencies in early modern human sites in Europe, but they are almost non-existent in Neanderthal sites (Hayden, 1990; Mellars, 1995). This difference could be taken to indicate that Neanderthals did not use clothing at all. However, there are two reasons to think that this interpretation is incorrect. First, use-wear studies indicate that Neanderthals employed other types of stone tools for hide processing (e.g. Beyries, 1988; Anderson-Gerfaud, 1990; Hayden, 1990). Hide processing made up a small percentage of the tasks identified in these studies, but the percentage was not negligible. In addition, Soressi et al. (2013) have recently reported the discovery of bone *lissoirs* that were used for hide processing at two Middle Palaeolithic sites in Europe. Together, these observations suggest that Neanderthals engaged in hide processing on a regular basis, but not with the same frequency as early modern humans. This, we contend, is the pattern predicted by the hypothesis that Neanderthals used cape-like clothing while early modern humans used specialized cold weather clothing. So, it appears that the hypothesis is actually supported by four different lines of evidence.

If the Neanderthals only had cape-like clothing while the early modern humans had specialized cold weather clothing, there are implications for the replacement of the Neanderthals by early

modern humans. As explained earlier, it is generally agreed that the Neanderthals' stout body and relatively short limbs are adaptations to glacial conditions (Feldesman et al., 1990; Ruff, 1993). In contrast, the bodies of early modern humans appear to have been adapted for tropical conditions (Ruff, 1993). Thus, the implications depend on the advantage of early modern human clothing over Neanderthal clothing compared to the advantage of Neanderthal physiology over early modern human physiology. If early modern human clothing conferred a small advantage relative to the advantage conferred by Neanderthal physiology, then early modern humans' clothing may have simply helped level the playing field in relation to Neanderthal physiology, allowing them to cope with cold temperatures as well as Neanderthals could. Alternatively, if the advantage conferred by early modern human clothing was large compared to the advantage conferred by Neanderthal physiology, then early modern humans' clothing could have helped them outcompete Neanderthals. There are at least two reasons for favouring the second of these scenarios. One is that energetic modeling work suggests the lower critical temperature and minimum sustainable ambient temperature for Neanderthals would have been only 1–2.5 °C below those for early modern humans (Aiello and Wheeler, 2003), which suggests that the advantages conferred by Neanderthal physiology were relatively small. The other is that there is evidence that early modern humans were able to withstand more adverse conditions than Neanderthals. Their cold period occupations extend further north than do those of Neanderthals (van Andel et al., 2003), and attempts to reconstruct climatic conditions at archaeological sites dating to OIS3 indicate that, on average, early modern human-associated sites experienced higher wind chill than Neanderthal-associated sites (Aiello and Wheeler, 2003). This implies that, with respect to cold stress, early modern humans' physiological disadvantages were outweighed by their technological advantages. There is reason therefore to believe that the difference in clothing helped early modern humans successfully compete with Neanderthals for territory and resources.

Why there was a difference in the clothing used by Neanderthals and early modern humans is unclear. There appear to be two possibilities. One is that the Neanderthals were insufficiently intelligent to create garments of the same thermal effectiveness as those used by early modern humans. Klein (2009) has argued that early modern humans were able to “wield culture more effectively” (p. 1526) than Neanderthals due to a change in their brain function ca. 50 kya. According to Klein, this change enhanced the ability of early modern humans to communicate symbolically and allowed them to produce tools from a wider range of materials. It is feasible that the manufacture of specialized cold weather clothing is another consequence of the putative change in brain function. The other possibility is that cultural factors explain the difference in clothing. Recent history demonstrates that differences in cognitive ability are not needed for one hominin population to expand at the expense of another, or for one hominin population to cope with a deteriorating climate while another does not—cultural factors are sufficient. Cultural innovations enabled early farmers to expand into areas of Europe already occupied by hunter-gatherer groups, and cultural factors explain why the Greenland Norse died out while the Inuit Greenlanders expanded their territory during the Little Ice Age (Henrich, 2010). Thus, it is feasible that the difference in clothing was a consequence of differences in cultural trajectories, combined with social and/or linguistic impediments to cultural transmission from early modern humans to Neanderthals. For example, it is possible that the Neanderthals' failure to develop more thermally effective clothing relates to the procurement of pelts. Trapping appears to be required for successful hunting of mustelids and canids (Holliday,

1998). Thus, it could be that Neanderthals failed to develop as thermally effective clothing as early modern humans because their trapping technology was less productive than that developed by early modern humans, as Holliday and Churchill (2006) have suggested. Determining which, if any, of these hypotheses is correct will require further empirical research.

### Acknowledgments

MC is grateful to Karen Lupo and Brian Coddling for the invitation to participate in the symposium honouring Jim O'Connell at the 2015 annual meeting of the Society for American Archaeology in San Francisco, and for the invitation to contribute to this special issue of the *Journal of Anthropological Archaeology*. We thank Conrad Brimacombe, Kate Britton, Keith Dobney, Mana Dembo, Marina Elliott, Ian Gilligan, Brian Hayden, Trenton Holliday, Vance Hutchinson, Steve Kuhn, Dana Lepofsky, Lee Lyman, Luseadra McKerracher, Kim Plomp, Bernard Wood, and an anonymous reviewer for their comments on earlier versions of this paper. Ian Gilligan's comments in particular resulted in major changes to the interpretation of the results. MC is supported by the Social Sciences and Humanities Research Council of Canada, the Canada Research Chairs Program, the Canada Foundation for Innovation, the British Columbia Knowledge Development Fund, and Simon Fraser University. LT's work on the study reported here was supported by the Social Sciences and Humanities Research Council of Canada (award no. 755-2011-0406) and Simon Fraser University – Canada. We thank all these funding bodies. Last but not least, MC would like to express his gratitude to Jim O'Connell for his friendship and guidance over nearly 20 years.

### Appendix A

Mammalian taxa represented in the Stage 3 Project Faunal Database.

Family	Taxon	Common name(s) (45)
Bovidae	<i>Bos primigenius</i>	Auroch
	<i>Bos</i> sp.	Auroch, cattle, kouprey, banteng, gaur, seladang, yak
	<i>Bison</i> sp./ <i>B. priscus</i>	Bison, wisent, buffalo
	<i>Capra</i> sp.	Goat, ibex, tur, markhor
	<i>Capra ibex</i>	Ibex
	<i>Capra pyrenaica</i>	Spanish ibex
	<i>Ovis</i> sp.	Sheep, mouflon, urial, argali
	<i>Ovibos moschatus</i>	Musk ox
	<i>Saiga tartarica</i>	Saiga antelope, saiga
	Undifferentiated bovids	
Canidae	<i>Canis lupus</i>	Wolf
	<i>Cuon</i> sp.	Dholes
	<i>Vulpes vulpes</i>	Red fox
	<i>Vulpes corsac</i>	Corsac fox
	<i>Alopex lagopus</i>	Arctic fox
Castoridae	<i>Castor fiber</i>	Beaver
Cercopithecidae	<i>Macaca sylvana</i> / <i>sylvanus</i>	Macaque, barbary ape

## Appendix A (continued)

Family	Taxon	Common name(s) (45)	
Cervidae	<i>Dama dama</i>	Fallow deer	
	<i>Megaloceros</i> sp./ <i>Megaloceros giganteus</i>	Giant deer	
	<i>Alces alces</i>	Elk, moose	
	<i>Cervus elaphus</i>	Red deer, wapiti, elk	
	<i>Rangifer tarandus</i>	Reindeer, caribou	
	<i>Capreolus capreolus</i>	Roe deer	
	<i>Rupicapra rupicapra</i>	Chamois	
	Dipodidae	<i>Sicista</i> sp.	Birch mice
		<i>Sicista subtilis</i>	Southern birch mouse
		<i>Sicista betulina</i>	Northern birch mouse
<i>Allactaga</i> sp.		Jerboas	
<i>Alactagulus</i> sp.		Jerboas, little earth hares	
Elephantidae	<i>Mammuthus primigenius</i>	Mammoth, woolly mammoth	
	<i>Elephas antiquus</i> / <i>Elephas (Palaeoloxodonta) sp.</i>	Straight-tusked elephants	
	Equidae	<i>Equus hydruntinus</i>	Steppe ass
		<i>Equus hemionus</i>	Hemione, kulan, onager
		<i>Equus caballus</i>	Wild horse
<i>Equus</i> sp.		Horse, kiang, zebra, quagga, ass, burro, donkey, hemione, kulan, onager	
	<i>Equus asinus</i>	Ass, burro, donkey	
	<i>Equus latipes</i>	Extinct horse	
Erinaceidae	<i>Erinaceus</i> sp.	Eurasian hedgehogs	
	<i>Erinaceus europea</i>	Hedgehog	
Felidae	<i>Panthera</i> sp.	Big cats	
	<i>Panthera leo</i>	Lion	
	<i>Panthera spealea</i>	Cave lion	
	<i>Panthera pardus</i>	Leopard	
	<i>Felis sylvestris</i>	Wild cat	
	<i>Lynx lynx</i> / <i>Felis lynx</i> / <i>Felis sp.</i>	Lynx	
	<i>Lynx</i> / <i>Felis pardina</i>	Pardel lynx, Spanish lynx	
	<i>Lynx</i> sp.	Undifferentiated lynxes	
Hyaenidae	<i>Crocuta crocuta</i> / <i>Hyaena spelaea</i>	Spotted hy(a)ena, striped hy(a)ena	
	Hystricidae	<i>Hystrix cristata</i> / <i>vinogradovi</i>	Porcupine, Old World porcupine
Leporidae		<i>Oryctolagus cuniculus</i>	Rabbit
	<i>Lepus</i> sp.	Hares, jack rabbits	
	<i>Lepus europaeus</i> / <i>capensis</i>	Brown hare	
	<i>Lepus timidus</i>	Mountain hare	
	<i>Lepus tolai</i>	Tolai hare	

## Appendix A (continued)

Family	Taxon	Common name(s) (45)
Muridae	<i>Apodemus</i> sp.	Old World mice
	<i>Apodemus flavicollis</i>	Yellow-necked mouse
	<i>Apodemus sylvaticus</i>	Wood mouse
	<i>Cricetus cricetus</i>	Common hamster
	<i>Cricetus</i> sp.	Hamsters
	<i>Cricetulus migratorius</i>	Grey hamster
	<i>Mesocricetus</i> sp.	Golden hamster
	<i>Allocricetus bursae</i>	Extinct hamster
	<i>Arvicola sapidus</i>	Southern water vole
	<i>Arvicola terrestris</i>	Northern water vole
	<i>Arvicola</i> sp.	Water vole
	<i>Pliomys lenki</i>	Extinct vole
	<i>Pliomys</i> sp.	Extinct voles
	<i>Ptericola (Pitymys) fatioi</i> (subgenus of <i>Microtus</i> )	No common name found
	<i>Ptericola (Pitymys) duodecimcostatus</i>	No common name found
	<i>Ptericola (Pitymys) subterraneus</i>	No common name found
	<i>Pericola (Pitymys) savii</i>	No common name found
	<i>Ptericola (Pitymys) sp.</i>	No common name found
	<i>Microtus breccensis</i> / <i>carbnera</i>	No common name found
	<i>Microtus afghanus</i>	No common name found
	<i>Microtus hyperboreus</i>	No common name found
	<i>Microtus arvalis</i>	Common vole, vole
	<i>Microtus agrestis</i>	Field vole
	<i>Microtus gregalis</i>	Vole, meadow mouse
	<i>Microtus nivalis</i>	Snow vole
	<i>Microtus oeconomus</i> / <i>raticeps</i>	Northern vole
	<i>Microtus</i> sp.	Undifferentiated voles, meadow mice
	<i>Clethrionomys glareolus</i>	Bank vole
	<i>Clethrionomys rutilus</i>	Ruddy vole
<i>Clethrionomys rufocanus</i>	Grey sided vole	
<i>Clethrionomys</i> sp.	Red backed mice	
<i>Lemmus</i> sp.	Lemmings	
<i>Lemmus lemmus</i>	Norway lemming	
<i>Dicrostonyx torquatus</i>	Collared lemming	
<i>Lagurus lagurus</i>	Steppe lemming	
<i>Eolagurus luteus</i>	No common name found	
<i>Myopus schisticolor</i>	Wood lemming	
<i>Spalax</i> sp.	Mole-rats	
<i>Ellobius</i> sp.	Mole-voles, mole-lemmings	

(continued on next page)

## Appendix A (continued)

Family	Taxon	Common name(s) (45)	
Mustelidae	<i>Meles meles</i>	Badger	
	<i>Mustela minuta</i>	Least weasel	
	<i>Mustela erminea</i>	Stoat, ermine	
	<i>Mustela nivalis</i>	Weasel	
	<i>Mustela</i> sp.	Mink, ferret, weasel, stoat, ermine, polecat	
	<i>Putorius/Mustela putorius</i>	Polecat	
	<i>Putorius putorius robusta/Mustela eversmanni</i>	Large polecat, steppe polecat	
	<i>Martes</i> sp.	Marten, fisher, sable	
	<i>Gulo gulo</i>	Wolverine	
	<i>Lutra lutra</i>	Otter	
	Myoxidae	<i>Myoxus/Glis glis</i>	Edible doremouse, fat doremouse
		<i>Muscardinus avellanarius</i>	Doremouse, hazel mouse
		<i>Eliomys quercinus</i>	Garden doremouse
Ochotonidae	<i>Ochotona pusilla</i>	Pika, mouse hare, coney	
Rhinocerotidae	<i>Dicerorhinus/Stephanorhinus mercki/kirckbergensis</i>	Extinct rhino	
	<i>Dicerorhinus/Stephanorhinus hemitoechus</i>	Extinct rhino	
	<i>Dicerorhinus/Stephanorhinus</i> sp.	Hairy rhino, Sumatran rhino	
	<i>Coelodonta antiquitatis</i>	Woolly rhino	
	Soricidae	<i>Crociodura</i> sp. group	White toothed shrews
<i>Sorex "araneus"</i> group		Common shrews	
<i>Sorex minutus</i>		Pigmy shrew	
<i>Sorex minutissimus</i>		Least shrew	
<i>Sorex</i> sp.		Long-tailed shrews	
<i>Neomys</i> sp.		Water shrews	
Sciuridae		<i>Pteromys</i> sp.	Flying squirrels
	<i>Marmota</i>	Extinct marmot	
	<i>primigenia</i>	No common name found	
	<i>Marmota bobak</i>	Marmot, woodchuck, groundhog	
	<i>Marmota marmota</i>	Alpine marmot	
	<i>Spermophilus major/Citellus superciliosus</i>	No common name found	
	<i>Spermophilus</i> sp.	Ground squirrels, susliks	
	<i>Sciurus vulgaris</i>	Red squirrel	
Suidae	<i>Sus scrofa</i>	Wild boar, wild pig, pig	

## Appendix A (continued)

Family	Taxon	Common name(s) (45)
Talpidae	<i>Talpa</i> sp.	Old World moles
	<i>Talpa europea</i>	Mole
	<i>Desmana</i> sp.	Desmans
Ursidae	<i>Ursus arctos</i>	Brown bear, grizzly, grizzly bear
	<i>Ursus</i> sp.	Bears
	<i>Ursus spelaea</i>	Cave bear
Vespertilionidae	<i>Plecotus auritus</i>	Old World long-eared bat

## References

- Adler, D.S., Wilkinson, K.N., Blockley, S., Mark, D.F., Pinhasi, R., Schmidt-Magee, B.A., Nahapetyan, S., Mallol, C., Berna, F., Glauberman, P.J., Raczyński-Henk, Y., Wales, N., Frahm, E., Jöris, O., MacLeod, A., Smith, V.C., Cullen, V.L., Gasparian, B., 2014. Early Levallois technology and the Lower to Middle Paleolithic transition in the Southern Caucasus. *Science* 345, 1609–1613.
- Aiello, L., Wheeler, P., 2003. Neanderthal thermoregulation and the glacial climate. In: van Andel, T., Davies, W. (Eds.), *Neanderthals and Modern Humans in the European Landscape during the Last Glaciation*. McDonald Institute for Archaeological Research, Cambridge, pp. 147–166.
- Anderson-Gerfaud, P., 1990. Aspects of behaviour in the Middle Palaeolithic: functional analysis of stone tools from Southwest-France. In: Mellars, P. (Ed.), *The Emergence of Modern Humans: An Archaeological Perspective*. Edinburgh University Press, Edinburgh, pp. 389–418.
- Backwell, L., d'Errico, F., Wadley, L., 2008. Middle stone age bone tools from the Howiesons Poort layers, Sibudu Cave, South Africa. *J. Archaeol. Sci.* 35, 1566–1580.
- Bar-Yosef, O., Bordes, J.G., 2010. Who were the makers of the Châtelperronian culture? *J. Hum. Evol.* 59, 586–593.
- Bemilli, C., Bayles, G., 2009. Aurignacian animal exploitation at Solutré (Saône-et-Loire, France). In: Fontana, L., Chauvière, F.-X., Bridault, A. (Eds.), *In Search of Total Animal Exploitation by Upper Palaeolithic and Mesolithic Hunter-Gatherer Societies*, Proceedings of the XVth UISPP Congress, Session C61. British Archaeological Reports, Oxford, pp. 37–47.
- Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. Roy. Stat. Soc. Ser. B: Stat. Methodol.* 57, 289–300.
- Beyries, S., 1988. Functional variability of lithic sets in the Middle Paleolithic. In: Dibble, H., Montet-White, A. (Eds.), *Upper Pleistocene Prehistory of Western Eurasia*. University of Pennsylvania Press, Philadelphia, pp. 213–223.
- Bogoraz-Tan, V.G., 1909. *The Chukchee: Material Culture, Religion, Social Organization*. Brill, New York.
- Bouchud, J., 1975. Étude de la faune de l'Abri Pataud. In: Movius, H.L. (Ed.), *Excavation of the Abri Pataud, Les Eyzies (Dordogne)*. Peabody Museum of Archaeology and Ethnology, Cambridge, MA, pp. 69–153.
- Boyle, K.V., 1998. The Middle Palaeolithic Geography of Southern France: Resources and Site Location. *British Archaeological Reports*, Oxford.
- Cartmill, M., Smith, F.H., 2009. *The Human Lineage*. Blackwell, Hoboken, NJ.
- Castel, J., 2011. Archéozoologie de l'Aurignacien de l'Abri Castanet (Sergeac, Dordogne, France): Les fouilles 1994–1998. *Rev. Paléobiol. Genève* 30, 783–815.
- Castel, J.-C., Discamps, E., Soulier, M.-C., Sandgathe, D.M., Dibble, H.L., McPherron, S. P., Goldberg, P., Turq, A., in press. Neandertal subsistence strategies during the Quina Mousterian at Roc de Marsal (France). *Quatern. Int.* <http://dx.doi.org/10.1016/j.quaint.2015.12.033>.
- Collard, M., Dembo, M., 2013. Modern human origins. In: Begun, D. (Ed.), *A Companion to Paleoanthropology*. Wiley, Malden, MA, pp. 557–581.
- Cotel, A.J., Golingo, R., Oakes, J.E., Riewe, R.R., 2004. Effect of ancient Inuit fur parka ruffs on facial heat transfer. *Clim. Res.* 26 (77–84), 31.
- Crépin, L., Péan, S., Láznicková-Galetová, M., 2014. Comportements de subsistance au Paléolithique supérieur en Crimée: Analyse archéozoologique des couches 6–2, 6–1 et 5–2 de Buran-Kaya III. *L'Anthropologie* 118, 584–598.
- Debénath, A., Jelínek, A.J., 1998. Nouvelles fouilles à la Quina (Charente): résultats préliminaires. *Gallia Préhist.* 40, 29–74.
- Dobrowolskaya, M., Richards, M., Trinkaus, E., 2012. Direct radiocarbon dates for the mid Upper Paleolithic (Eastern Gravettian) burials from Sungir, Russia. *Bull. Mém. Soc. Anthropol. Paris* 24, 96–102.
- Douka, K., Higham, T.F., Wood, R., Boscatto, P., Gambassini, P., Karkanas, P., Peresani, M., Ronchitelli, A.M., 2014. On the chronology of the Uluzzian. *J. Hum. Evol.* 68, 1–13.
- Feldesman, M.R., Kleckner, J.G., Lundy, J.K., 1990. Femur/stature ratio and estimates of stature in mid-and late-Pleistocene fossil hominids. *Am. J. Phys. Anthropol.* 83, 359–372.

- Finlayson, C., 2004. Neanderthals and Modern Humans: An Ecological and Evolutionary Perspective. Cambridge University Press, Cambridge.
- Finlayson, C., 2009. The Humans Who Went Extinct: Why Neanderthals Died Out and We Survived. Oxford University Press, New York.
- Froehle, A.W., Churchill, S.E., 2009. Energetic competition between Neandertals and anatomically modern humans. *PaleoAnthropology* 2009, 96–116.
- Fu, Q., Li, H., Moorjani, P., Jay, F., Slepchenko, S.M., Bondarev, A.A., Johnson, P.L.F., Aximu-Petri, A., Prüfer, K., de Filippo, C., Meyer, M., Zwyns, N., Salazar-García, D. C., Kuzmin, Y.V., Keates, S.G., Kosintsev, P.A., Razhev, D.I., Richards, M.P., Peristov, N.V., Lachmann, M., Douka, K., Higham, T.F.G., Slatkin, M., Hublin, J.J., Reich, D., Kelso, J., Viola, F.B., Pääbo, S., 2014. Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature* 514, 445–449.
- Gautier, A., 1979. Documentation paléontologique. In: Hustaerts, P., de Heuzelen, J. (Eds.), *Le Site Paléolithique De Maisières-Canal. De Tempel, Brugge*, pp. 67–68.
- Gilligan, I., 2007. Neanderthal extinction and modern human behaviour: the role of climate change and clothing. *World Archaeol.* 39, 499–514.
- Golovanova, L.V., Doronichev, V.B., Cleghorn, N.E., 2010a. The emergence of bone-working and ornamental art in the Caucasian Upper Palaeolithic. *Antiquity* 84, 299–320.
- Golovanova, L.V., Doronichev, V.B., Cleghorn, N.E., Koulikova, M.A., Sapelko, T.V., Shackley, M.S., 2010b. Significance of ecological factors in the Middle to Upper Palaeolithic transition. *Curr. Anthropol.* 51, 655–691.
- Green, R.E., Krause, J., Briggs, A.W., Maricic, T., Stenzel, U., Kircher, M., Patterson, N., Li, H., Zhai, W., Fritz, M.H.-Y., Hansen, N.F., Durand, E.Y., Malaspinas, A.-S., Jensen, J.D., Marques-Bonet, T., Alkan, C., Prüfer, K., Meyer, M., Burbano, H.A., Good, J.M., Schultz, R., Aximu-Petri, A., Buffhof, A., Höber, B., Höffner, B., Siegemund, M., Weihmann, A., Nusbaum, C., Lander, E.S., Russ, C., Novod, N., Affourtit, J., Egholm, M., Verna, C., Rudan, P., Brajkovic, D., Kucan, Z., Gušić, I., Doronichev, V.B., Golovanova, L.V., Lalueza-Fox, C., de la Rasilla, M., Fortea, J., Rosas, A., Schmitz, R.W., Johnson, P.L.F., Eichler, E.E., Falush, D., Birney, E., Mullikin, J.C., Slatkin, M., Nielsen, R., Kelso, J., Lachmann, M., Reich, D., Pääbo, S., 2010. A draft sequence of the Neandertal genome. *Science* 328, 710–722.
- Hayden, B., 1990. The right rub: hide working in high ranking households. In: Gräslund, B., Knutsson, H., Knutsson, K., Taffinder, J. (Eds.), *The Interpretive Possibilities of Microwear Studies. Societas Archaeologica Upsaliensis*, Uppsala, pp. 89–102.
- Hayden, B., 1993. The cultural capacities of the Neanderthals: a review and re-evaluation. *J. Hum. Evol.* 24, 113–146.
- Henrich, J., 2010. The evolution of innovation-enhancing institutions. In: O'Brien, M. J., Shennan, S. (Eds.), *Innovation in Cultural Systems: Contributions from Evolutionary Anthropology*. MIT Press, Cambridge, MA, pp. 99–120.
- Higham, T., Douka, K., Wood, R., Ramsey, C.B., Brock, F., Basell, L., Camps, M., Arrizabalaga, A., Baena, J., Barroso-Ruiz, C., Bergman, C., Boitard, C., Boscato, P., Caparrós, M., Conard, N.J., Draily, C., Froment, A., Galván, B., Gambassini, P., García-Moreno, A., Grimaldi, S., Haesaerts, P., Holt, B., Iriarte-Chiapusso, M.J., Jelinek, A., Jordá Pardo, J.F., Maíllo-Fernández, J.M., Marom, A., Maroto, J., Menéndez, M., Metz, L., Morin, E., Moroni, A., Negrino, F., Panagopoulou, E., Peresani, M., Pirson, S., De La Rasilla, M., Riel-Salvatore, J., Ronchitelli, A., Santamaria, D., Semal, P., Slimak, L., Soler, J., Soler, N., Villaluenga, A., Pinhasi, R., Jacobi, R., 2014. The timing and spatiotemporal patterning of Neanderthal disappearance. *Nature* 512, 306–309.
- Hoffecker, J.F., 2005a. Innovation and technological knowledge in the Upper Paleolithic of northern Eurasia. *Evol. Anthropol.* 14, 186–198.
- Hoffecker, J.F., 2005b. A Prehistory of the North: Human Settlement of the Higher Latitudes. Rutgers University Press, New Brunswick, NJ.
- Holliday, T.W., 1997. Postcranial evidence of cold adaptation in European Neandertals. *Am. J. Phys. Anthropol.* 104, 245–258.
- Holliday, T.W., 1998. The ecological context of trapping among recent hunter-gatherers: implications for subsistence in terminal Pleistocene Europe. *Curr. Anthropol.* 39, 711–719.
- Holliday, T.W., Churchill, S.E., 2006. Mustelid hunting by recent foragers and the detection of trapping in the European Paleolithic. In: Haws, J.A., Hockett, B.S., Brugal, J.-P. (Eds.), *Paleolithic Zooarchaeology in Practice. Archaeopress*, Oxford, pp. 45–56.
- Itkonen, T.L., 1962. The Lapps in Finland up to 1945. Porvoo, Helsinki.
- Jefferies, R.W., 1990. A technological and functional analysis of Middle Archaic hafted endscrapers from the Black Earth site, Saline County, Illinois. *Midcont. J. Archaeol.* 15, 3–36.
- Jiménez-Espejo, F.J., Martínez-Ruiz, F., Finlayson, C., Paytan, A., Sakamoto, T., Ortega-Huertas, M., Finlayson, G., Iijima, K., Gallego-Torres, D., Fa, D., 2007. Climate forcing and Neanderthal extinction in southern Iberia: insights from a multiproxy marine record. *Quatern. Sci. Rev.* 26, 836–852.
- Kitagawa, K., Krönneck, P., Conard, N.J., Münzel, S.C., 2012. Exploring cave use and exploitation among cave bears, carnivores and hominins in the Swabian Jura, Germany. *J. Taphon.* 10, 439–461.
- Kittler, R., Kayser, M., Stoneking, M., 2003. Molecular evolution of *Pediculus humanus* and the origin of clothing. *Curr. Biol.* 13, 1414–1417.
- Kittler, R., Kayser, M., Stoneking, M., 2004. Erratum: molecular evolution of *Pediculus humanus* and the origin of clothing. *Curr. Biol.* 14, 2309.
- Klein, R.G., 2003. Whither the Neanderthals? *Science* 299, 1525.
- Klein, R.G., 2009. Darwin and the recent African origin of modern humans. *Proc. Natl. Acad. Sci. U.S.A.* 106, 16007–16009.
- Krause, J., Orlando, L., Serre, D., Viola, B., Prüfer, K., Richards, M.P., Hublin, J.-J., Hänni, C., Derevianko, A.P., Pääbo, S., 2007. Neanderthals in central Asia and Siberia. *Nature* 449, 902–904.
- Laquay, G., 1981. Recherches sur les faunes du Würm I en Périgord. Unpublished PhD Dissertation. Université de Bordeaux I.
- Loebel, T.J., 2013. Endscrapers, use-wear, and early Paleoindians in Eastern North America. In: Gingerich, J.A.M. (Ed.), *The Eastern Fluted Point Tradition*. University of Utah Press, Utah, pp. 315–330.
- Mellars, P.A., 1995. The Neanderthal Legacy: An Archaeological Perspective from Western Europe. Princeton University Press, Princeton, NJ.
- Mellars, P., 2004. Neanderthals and the modern human colonization of Europe. *Nature* 432, 461–465.
- Miracle, P., 2005. Late Mousterian subsistence and cave use in Dalmatia: the zooarchaeology of Mujina Pecina, Croatia. *Int. J. Osteoarchaeol.* 15, 84–105.
- Mizell, L.R., Frishman, D., Saxon, R., 1956. Development of Non-Critical Substitutes for Wolverine Fur. Textile Series Report 92.
- Montana Trappers Association, no date. Wolverine. <<http://www.montanatrappers.org/furbearers/wolverine.htm#>> (accessed 17-Dec-15).
- Morin, E., 2012. Reassessing Paleolithic Subsistence: The Neandertal and Modern Human Foragers of Saint-Césaire. Cambridge University Press, Cambridge.
- Mourre, V., Costamagno, S., Thiébaud, C., Allard, M., Bruxelles, L., Colonge, D., Cravinho, S., Jeannot, M., Juillard, F., Laroulandie, V., Maureille, B., 2008. Le site moustérien de la Grotte du Noisetier à Fréchet-Aure (Hautes-Pyrénées) – premiers résultats des nouvelles fouilles. In: Jaubert, J., Bordes, J.-G., Ortega, I. (Eds.), *Les Sociétés Paléolithiques D'un Grand Sud-Ouest: Nouveaux Gisements, Nouvelles Méthodes, Nouveaux Résultats – Actes Des Journées Décentralisées De La SPF Des 24–25 Novembre 2006. Société Préhistorique Française, Nanterre, France*, pp. 189–202.
- Musil, R., 2003. The Middle and Upper Palaeolithic game suite in central and southeastern Europe. In: van Andel, T., Davies, W. (Eds.), *Neanderthals and Modern Humans in the European Landscape during the Last Glaciation: Archaeological Results of the Stage 3 Project*. McDonald Institute for Archaeological Research, Cambridge, pp. 167–190.
- Niven, L., 2013. A diachronic evaluation of Neanderthal cervid exploitation and site use at Pech de l'Azé IV, France. In: Clark, J.L., Speth, J.D. (Eds.), *Zooarchaeology and Modern Human Origins: Human Hunting Behavior during the Later Pleistocene*. Springer, Dordrecht; The Netherlands, pp. 151–161.
- Niven, L., Steele, T.E., Rendu, W., Mallye, J.B., McPherron, S.P., Soressi, M., Jaubert, J., Hublin, J.J., 2012. Neanderthal mobility and large-game hunting: the exploitation of reindeer during the Quina Mousterian at Chez-Pinaud Jonzac (Charente-Maritime, France). *J. Hum. Evol.* 63, 624–635.
- Nowak, R.M., 1999. Walker's Mammals of the World. John Hopkins University Press, Baltimore, MD.
- Oakes, J., Wilkins, H., Riewe, R., Kelker, D., Forest, T., 1995. Comparison of traditional and manufactured cold weather ensembles. *Clim. Res.* 5, 83–90.
- O'Connell, J.F., 2006. How did modern humans displace Neanderthals? Insights from hunter-gatherer ethnography and archaeology. In: Conard, N.J. (Ed.), *When Neanderthals and Modern Humans Met*. Kerns Verlag, Tübingen, pp. 43–64.
- Pryde, D., 1972. Nunaga: My Land, My Country. Hurtig, Edmonton, AB.
- Richards, M.P., Trinkaus, E., 2009. Isotopic evidence for the diets of European Neanderthals and early modern humans. *Proc. Natl. Acad. Sci. U.S.A.* 106, 16034–16039.
- Richter, J., 2011. When did the Middle Paleolithic begin? In: Conard, N.J., Richter, J. (Eds.), *Neanderthal Lifeways, Subsistence and Technology: One Hundred Fifty Years of Neanderthal Study*. Springer, Dordrecht, The Netherlands, pp. 7–14.
- Ruff, C.B., 1993. Climatic adaptation and hominid evolution: the thermoregulatory imperative. *Evol. Anthropol.* 2, 53–60.
- Ruff, C.B., Trinkaus, E., Holliday, T.W., 1997. Body mass and encephalization in Pleistocene *Homo*. *Nature* 387, 173–176.
- Sandgathe, D.M., Dibble, H.L., Goldberg, P., McPherron, S.P., Turq, A., Niven, L., Hodgkins, J., 2011. On the role of fire in Neanderthal adaptations in Western Europe: evidence from Pech de l'Azé IV and Roc de Marsal, France. *PaleoAnthropology* 2011, 216–242.
- Schreie, D.C., 2006. The taphonomy of a Middle Devensian (MIS 3) vertebrate assemblage from Lynford, Norfolk, UK, and its implications for Middle Paleolithic subsistence strategies. *J. Quatern. Sci.* 21, 543–556.
- Shott, M.J., Weedman, K.J., 2007. Measuring reduction in stone tools: an ethnoarchaeological study of Gamo hide scrapers from Ethiopia. *J. Archaeol. Sci.* 34, 1016–1035.
- Sørensen, B., 2009. Energy use by Eem Neanderthals. *J. Archaeol. Sci.* 36, 2201–2205.
- Soressi, M., Rendu, W., Texier, J.-P., Daulny, É.C.L., D'Errico, F., Laroulandie, V., Maureille, B., Niclot, M., Schwartz, S., Tillier, A., 2008. Pech-de-l'Azé I (Dordogne, France): nouveau regard sur un gisement moustérien de tradition acheuléenne connu depuis le XIXe siècle. In: Jaubert, J., Bordes, J.-G., Ortega, I. (Eds.), *Les Sociétés Paléolithiques D'un Grand Sud-Ouest: Nouveaux Gisements, Nouvelles Méthodes, Nouveaux Résultats – Actes Des Journées Décentralisées De La SPF Des 24–25 Novembre 2006. Société Préhistorique Française, Nanterre, France*, pp. 95–132.
- Soressi, M., McPherron, M.P., Lenoir, M., Dogandžić, T., Goldberg, P., Jacobs, Z., Maigrot, Y., Martisius, N.L., Miller, C.E., Rendu, W., Richards, M., Skinner, M.M., Steele, T.E., Talamo, S., Texier, J.-P., 2013. Neanderthals made the first specialized bone tools in Europe. *Proc. Natl. Acad. Sci. U.S.A.* 110, 14186–14190.
- Soulier, M., Mallye, J., 2012. Hominid subsistence strategies in the south-west of France: a new look at the early Upper Palaeolithic faunal material from Roc-de-Combe (Lot, France). *Quatern. Int.* 252, 99–108.
- Stanford, Q.H., 2003. Canadian Oxford World Atlas. Oxford University Press, Toronto.
- Stenton, D.R., 1991. The adaptive significance of caribou winter clothing for arctic hunter-gatherers. *Etud. Inuit Stud.* 15, 3–28.

- Stewart, J.R., 2007. Neanderthal extinction as part of the faunal change in Europe during oxygen isotope stage 3. *Acta Zool. Cracov.* 50, 93–124.
- Stewart, J.R., van Kolfschoten, T., Markova, A., Musil, R., 2003. The mammalian faunas of Europe during oxygen isotope stage three. In: Van Andel, T.H., Davies, W. (Eds.), *Neanderthals and Modern Humans in the European Landscape during the Last Glaciation*. McDonald Institute for Archaeological Research, Cambridge, pp. 103–130.
- Stiner, M.C., 2001. Thirty years on the “Broad spectrum revolution” and Paleolithic demography. *Proc. Natl. Acad. Sci. U.S.A.* 98, 6993–6996.
- Stiner, M.C., Kuhn, S.L., 2006. Changes in the ‘connectedness’ and resilience of Paleolithic societies in Mediterranean ecosystems. *Hum. Ecol.* 34, 693–712.
- Stiner, M.C., Barkai, R., Gopher, A., 2009. Cooperative hunting and meat sharing 400–200 kya at Qesem Cave, Israel. *Proc. Natl. Acad. Sci. U.S.A.* 106, 13207–13212.
- Stringer, C., 2002. Modern human origins: progress and prospects. *Philos. Trans. Roy. Soc. Lond. B: Biol. Sci.* 357, 563–579.
- Stringer, C., Palike, H., Van Andel, T.H., Huntley, B., Valdes, P.J., Allen, J.R.M., 2003. Climatic stress and the extinction of the Neanderthals. In: van Andel, T.H., Davies, W. (Eds.), *Neanderthals and Modern Humans in the European Landscape during the Last Glaciation*. McDonald Institute for Archaeological Research, Cambridge, pp. 233–240.
- Stringer, C.B., 2006. The Neanderthal-*H. sapiens* interface in Eurasia. In: Harvati, K., Harrison, T. (Eds.), *Neanderthals Revisited: New Approaches and Perspectives*. Springer, Dordrecht, The Netherlands, pp. 315–323.
- Svoboda, J., Péan, S., Wojtal, P., 2005. Mammoth bone deposits and subsistence practices during Mid-Upper Palaeolithic in Central Europe: three cases from Moravia and Poland. *Quatern. Int.* 126, 209–221.
- Toups, M.A., Kitchen, A., Light, J.E., Reed, D.L., 2011. Origin of clothing lice indicates early clothing use by anatomically modern humans in Africa. *Mol. Biol. Evol.* 28, 29–32.
- van Andel, T.H., 2003. Glacial environments I: the Weichselian climate in Europe between the end of the OIS-5 interglacial and the Last Glacial Maximum. In: van Andel, T.H., Davies, W. (Eds.), *Neanderthals and Modern Humans in the European Landscape during the Last Glaciation*. McDonald Institute for Archaeological Research, Cambridge, pp. 9–19.
- van Andel, T.H., Davies, W., 2003. Neanderthals and Modern Humans in the European Landscape during the Last Glaciation. McDonald Institute for Archaeological Research, Cambridge.
- van Andel, T., Davies, W., Weninger, B., 2003. The human presence in Europe during the Last Glacial period I: human migrations and the changing climate. In: van Andel, T.H., Davies, W. (Eds.), *Neanderthals and Modern Humans in the European Landscape during the Last Glaciation*. McDonald Institute for Archaeological Research, Cambridge, pp. 31–56.
- Wales, N., 2012. Modeling Neanderthal clothing using ethnographic analogues. *J. Hum. Evol.* 63, 781–795.
- White, M.J., 2006. Things to do in Doggerland when you’re dead: surviving OIS3 at the northwestern-most fringe of Middle Palaeolithic Europe. *World Archaeol.* 38, 547–575.
- Wolpoff, M.H., Hawks, J., Caspari, R., 2000. Multiregional, not multiple origins. *Am. J. Phys. Anthropol.* 112, 129–136.
- Wolpoff, M., Mannheim, B., Mann, A., Hawks, J., Caspari, R., Rosenberg, K., Frayer, D., Gill, G., Clark, G., 2004. Why not the Neandertals? *World Archaeol.* 36, 527–546.